

# Effects of Non-Commutativity on Light Hydrogen-Like Atoms and Proton Radius

M. Moumni

Department of Matter Sciences, University of Biskra; Algeria  
m.moumni@yahoo.fr

A. BenSlama

Department of Physics, University Constantine1; Algeria  
a.benslama@yahoo.fr

November 1, 2013

## Abstract

We study the corrections induced by the theory of non-commutativity, in both space-space and space-time versions, on the spectrum of hydrogen-like atoms. For this, we use the relativistic theory of two-particle systems to take into account the effects of the reduced mass, and we use perturbation methods to study the effects of non-commutativity. We apply our study to the muon hydrogen with the aim to solve the puzzle of proton radius [R. Pohl et al., Nature 466, 213 (2010) and A. Antognini et al., Science 339, 417 (2013)]. The shifts in the spectrum are found more noticeable in muon  $H$  ( $\mu H$ ) than in electron  $H$  ( $eH$ ) because the corrections depend on the mass to the third power; This explains the discrepancy between  $\mu H$  and  $eH$  results. In space-space non-commutativity, the parameter required to resolve the puzzle  $\theta_{ss} \approx (0.35 GeV)^{-2}$ , exceeds the limit obtained for this parameter from various studies on  $eH$  Lamb shift. For space-time non-commutativity, the value  $\theta_{st} \approx (14.3 GeV)^{-2}$  has been obtained and it is in agreement with the limit determined by Lamb shift spectroscopy in  $eH$ . We have also found that this value fills the gap between theory and experiment in the case of  $\mu D$  and improves the agreement between theoretical and experimental values in the case of hydrogen-deuterium isotope shift.

**Keywords:** Non-Commutativity; H-Like Atoms; Proton Radius

**PACS:** 02.40.Gh, 67.63.Gh, 31.30.jr

## 1 Introduction

Historically, experimental spectroscopy was the perfect test for any theory having any connection with matter and it played the leading role in calibrating the values of physical constants. But lately, it has reached such precision that it

accessed the role of indicator of new theories; and last experience on the Lamb shift in muonic hydrogen [1] is the perfect example.

The discrepancy between these experimental results  $r_{\mu H} = 0.84169(66)fm$  and those extracted from electronic hydrogen or elastic electron-proton scattering and recorded in CODATA  $r_{eH} = 0.8775(51)fm$  [2] (The values are at  $7\sigma$  variance with respect to each other) has had an impact in the whole scientific community and this raised many questions about the cause of such disagreement.

The experimental methods used to obtain the two results are very elaborated. This is why many studies have investigated how to explain this difference and to reconcile the two results by trying to rectify the theory. But the difference between the two experiments remains a puzzle until now and especially after being reinforced recently with a more accurate value  $r_{\mu H} = 0.84087(39)$  [3].

Non-perturbative numerical computations of the Dirac equation confirmed the validity of perturbation methods used to compute the radius [4]-[5]. No significant QED correction has been found yet, which would explain the discrepancy [6]-[7]. Using electron scattering experiments, [8] found that data rules out values of the third Zemach moment large enough to explain the puzzle. Three-body physics does not solve the problem as demonstrated in [9]. Constraints from low energy data disfavor new spin-0, spin-1 and spin-2 particles as an explanation [10]. There are some claims that proton polarizability contribution in the Lamb shift may explain the discrepancy because it is proportional to the lepton mass to the fourth power [11]-[12]. These effects could be probed in scattering experiment planned to run at Paul Scherrer Institute (PSI). For more information about the different approaches to the problem, see references [6]-[13]-[14]-[15].

Because Pohl et al. used an indirect method to calculate  $r_{\mu H}$  that involves comparing the frequency measured experimentally with that given theoretically according to the radius [1]:

$$\frac{\Delta(2P_{1/2} \rightarrow 2S_{1/2})}{meV} = 206.0573(45) - 5.2262 \frac{r_p^2}{fm^2} + 0.0347 \frac{r_p^3}{fm^3} \quad (1a)$$

$$\frac{\Delta(2P_{3/2} \rightarrow 2S_{1/2})}{meV} = 209.9779(49) - 5.2262 \frac{r_p^2}{fm^2} + 0.0347 \frac{r_p^3}{fm^3} \quad (1b)$$

we propose, in this work, to modify the precedent theoretical expressions of the transition frequency by incorporating the corrections induced by the non-commutative structure of space-time.

The idea of taking non-commutative space-time coordinates dates from the thirties of last century. It had as objective to avoid infinities in Coulomb potentials (gravitation & electricity) by introducing an lower bound for the measurement of length. Despite the fact that the concept was suffering from some problems with unitarity and causality, the theory evolved from the mathematical point of view, especially after the work of Connes in the eighties of last century [16].

In 1999, the work of Seiberg and Witten on string theory [17] has aroused

new interest in the theory. They showed that the dynamics of the endpoints of an open string on a D-brane in the presence of a magnetic back-ground field is described by a theory of Yang-Mills on a non-commutative space-time.

Today, we find non-commutativity in various fields of physics such as solid state physics, where it was shown that is the framework in which Hall conductivity is quantized [18] and that it is the proper tool replacing Bloch's theory whenever translation invariance is broken in aperiodic solids [19]. In fluid mechanics, non-commutative fluids are introduced by studying the quantum Hall effect [20] or bosonization of collective fermion states [21]. There is also some connection with quantum statistical physics [22], and it is also an interpretation of Ising-type models [23]. One can even find a manifestation of the non-commutativity in the physiology of the brain, where non-commutative computation in the vestibulo-ocular reflex was demonstrated in a way that is unattainable by any commutative system [24]. [25] is an excellent reference for the different manifestations and applications of non-commutative field theory.

The theory is a distortion of space-time where the coordinates  $x^\mu$  become Hermitian operators and thus do not commute:

$$[x_{nc}^\mu, x_{nc}^\nu] = i\theta^{\mu\nu}; \mu, \nu = 0, 1, 2, 3 \quad (2)$$

The  $nc$  indices denote non-commutative coordinates.  $\theta^{\mu\nu}$  is the parameter of the deformation and it is an anti-symmetric real matrix. We distinguish two types of non-commutativity; the first one is the space-space case, where the deformation is introduced between the spatial coordinates only, and the second is when the spatial coordinates commute with one another but not with time coordinate and it is noted space-time case. For a review, one can see reference [26].

In the literature, there are a lot of studies on hydrogen atom in non-commutativity. For space-space non-commutativity, we cite [27] [28] [29] [30]. For the space-time case, one can see [31] [32] [33].

We have found in [31] and [32], that the corrections induced by non-commutativity on the spectrum of the hydrogen atom are proportional to the lepton mass to the third power (the result is confirmed by [33]), and this is exactly the shape of the corrections induced by the nuclear size as demonstrated in [34] [35]. We will apply our result to the muonic hydrogen, and we will incorporate therein the effects of the finite mass of the nucleus. We start by computing the corrections to the energies in both space-space and space-time cases of non-commutativity using perturbation methods in the Dirac theory of two particles systems. Then we compare to the difference between theoretical and experimental results obtained in  $\mu H$  experience. This allows us to obtain values of the non-commutative parameter that resolve the puzzle. Then we will discuss the possible effects of these corrections on the Lamb shift of muonic deuterium  $\mu D$  and on the difference between the radii of the proton and deuteron via the  $2S - 1S$  transition.

## 2 Coulomb Potential in Non-Commutative Space-Time

We start by rewriting (2) for the two versions to consider of the non-commutativity:

$$[x_{st}^j, x_{st}^0] = i\theta^{j0} \quad (3a)$$

$$[x_{ss}^j, x_{ss}^k] = i\theta^{jk} \quad (3b)$$

$st$  subscripts are for space-time case and  $ss$  ones are for space-space case. The 0 denotes time and Latin indices are used for space coordinates. To solve these relations, we follow [27] [28] [31] [32] and choose the Bopp shift formulation of the solutions [36]; we write:

$$x_{st}^j = x^j - i\theta^{j0}\partial_0 \quad (4a)$$

$$x_{ss}^j = x^j - \frac{i}{2\hbar}\theta^{jk}\partial_k \quad (4b)$$

The usual coordinates of space  $x^j$  satisfy the usual canonical permutation relations and time  $x^0$  is unchanged in both cases.

We are dealing with the stationary quantum equations, and this allows us to consider the energy as a constant parameter. The kinetic energy does not change since it depends on the momentum that remains unchanged, thus we take the Coulomb potential and construct its non-commutative image. To achieve this, we have to write the expression of  $r_{nc}^{-1}$ , where  $nc$  denotes the two cases considered here:

$$\frac{1}{r_{nc}} = (\|\vec{r} + \vec{\vartheta}_{nc}\|)^{-1} = (\sum x_{nc}^j \cdot x_{nc}^j)^{-1/2} \quad (5)$$

$\vec{\vartheta}_{nc}$  is the non-commutative correction of position vector. We make the development in series and because of the smallness of the non-commutative parameter, as one can see from the bounds given in the literature [26], we restrict ourselves to the 1st order in  $\theta$  and neglect the higher order terms [27] [31]:

$$r_{st}^{-1} = \left( 1 + i \frac{\partial_0 \vec{r} \cdot \vec{\theta}_{st}}{r^3} + O(\theta^2) \right) \quad (6a)$$

$$r_{ss}^{-1} = \left( 1 + \frac{i}{2} \frac{(\vec{r} \times \vec{\partial}) \cdot \vec{\theta}_{ss}}{r^3} + O(\theta^2) \right) \quad (6b)$$

We have used the vectorial notation:

$$\vec{\theta}_{st} \equiv (\theta_{st}^1, \theta_{st}^2, \theta_{st}^3); \theta_{st}^j = \theta^{j0} \quad (7a)$$

$$\vec{\theta}_{ss} \equiv (\theta_{ss}^1, \theta_{ss}^2, \theta_{ss}^3); \theta^{jk} = \frac{1}{2}\varepsilon^{jkl}\theta_{ss}^l \quad (7b)$$

$$\vec{\partial} \equiv (\partial_1, \partial_2, \partial_3) \quad (7c)$$

Thus, one can write the non-commutative Coulomb potential (up to the 1st order  $\theta$ ) as follows:

$$V_{st}(r) = -\frac{Ze^2}{r} - \frac{Ze^2 E}{\hbar} \frac{\vec{r} \cdot \vec{\theta}_{st}}{r^3} + O(\theta_{st}^2) \quad (8a)$$

$$V_{ss}(r) = -\frac{Ze^2}{r} - \frac{Ze^2}{4\hbar} \frac{\vec{L} \cdot \vec{\theta}_{ss}}{r^3} + O(\theta_{ss}^2) \quad (8b)$$

where we have used the fact that  $i\partial_0\psi = H\psi = (E/\hbar)\psi$  and  $\vec{r}' \times i\hbar\vec{\partial}' = \vec{r}' \times \vec{p}' = \vec{L}$  the orbital momentum.

An adequate choice of the parameters is  $\vec{\theta}_{nc} = \theta^{r0}\vec{r}'/r = \theta_{nc}\vec{r}'/r$ . The writing is similar to that in [37] for space-space non-commutativity and in [31] and [38] for space-time case. The choice made in this paper allows us to write the non-commutative Coulomb potential as:

$$V_{st}(r) = -\frac{Ze^2}{r} - \frac{Ze^2 E \theta_{st}}{\hbar} \frac{1}{r^2} + O(\theta_{st}^2) \quad (9a)$$

$$V_{ss}(r) = -\frac{Ze^2}{r} - \frac{Ze^2(\vec{L} \cdot \vec{\theta}_{ss})}{4\hbar} \frac{1}{r^3} + O(\theta_{ss}^2) \quad (9b)$$

It was indicated in [31] that the effect of space-time noncommutativity (9a) is similar to the effect of an electric field of a radial dipole centered on the proton. Similarly, in [30], the effect of space-space noncommutativity (9b) was presented as equivalent to the effects of magnetic field or spin.

Now we can compute the corrections induced by this additional term using perturbative methods in both versions of non-commutativity.

### 3 Corrections of the Dirac Energies

We write the Dirac equation ( $\alpha_i = \gamma_0\gamma_i$  and  $\gamma_\mu$  are the Dirac matrices):

$$i\hbar\partial_0 = H\psi = (\vec{\alpha} \cdot \vec{p}) + m\gamma^0 + eA_0 \quad (10)$$

After coordinates deformation, we employ the standard Dirac equation but with the non-commutative Coulomb potential  $A_0^{(nc)} = -Ze/r_{nc}$ , so we get:

$$eA_0^{(st)} = -\frac{Ze^2}{r_{st}} = -\frac{Ze^2}{r} - \frac{Ze^2 E \theta_{st}}{\hbar} \frac{1}{r^2} + O(\theta_{st}^2) \quad (11a)$$

$$eA_0^{(ss)} = -\frac{Ze^2}{r_{ss}} = -\frac{Ze^2}{r} - \frac{Ze^2(\vec{L} \cdot \vec{\theta}_{ss})}{4\hbar} \frac{1}{r^3} + O(\theta_{ss}^2) \quad (11b)$$

As mentioned before, we restrict ourselves to the 1st order in  $\theta$ . The Hamiltonian can now be expressed as:

$$H = H_0 + H_{nc} = (\vec{\alpha} \cdot \vec{p}) + m\gamma^0 - Ze^2/r + \Delta H_{nc} \quad (12)$$

$\Delta H_{nc}$  is the non-commutative correction to the usual Dirac Hamiltonian  $H_0$ :

$$\Delta H_{st} = -Ze^2 (E/\hbar) \theta_{st} r^{-2} \quad (13a)$$

$$\Delta H_{ss} = -Ze^2 (\vec{L} \cdot \vec{\theta}_{ss}/4\hbar) r^{-3} \quad (13b)$$

The smallness of the parameter  $\theta$  allows us to consider noncommutative corrections with perturbation theory; to the 1st order in  $\theta$ , the corrections of the eigenvalues are:

$$\Delta E_{nc} = \langle \Delta H_{nc} \rangle = \langle \Psi(\vec{r}') | \Delta H_{nc} | \Psi(\vec{r}') \rangle \quad (14)$$

where the  $\Psi(\vec{r}')$  are the eigenstates of the Dirac Hamiltonian for Coulomb potential.

Because the space-time correction is central while space-space one is not (this is due to the presence of the  $\vec{L}$  operator in  $\Delta H_{ss}$  (13b)), we will treat the two cases separately.

### 3.1 1st Order Corrections in Space-Time Non-Commutativity

To compute the corrections (14) with  $\Delta H_{nc}$  given by (13a) and because all the parameters are constants except the coordinate  $r$ , we have:

$$\Delta E_{st} = \langle \Delta H_{st} \rangle = -Ze^2 (E/\hbar) \theta_{st} \langle \Psi(\vec{r}') | r^{-2} | \Psi(\vec{r}') \rangle \quad (15)$$

We can use the expression of the  $\Psi(\vec{r}')$  from the literature as done in [30], or employ the recurrence relations given in [39] as we have done in [31]. The obtained mean value for  $r^{-2}$  is:

$$\left\langle \frac{1}{r^2} \right\rangle = \frac{2\kappa(2\kappa\varepsilon - 1)(1 - \varepsilon^2)^{3/2}}{Z\alpha\sqrt{\kappa^2 - Z^2\alpha^2} [4(\kappa^2 - Z^2\alpha^2) - 1]} \left( \frac{mc}{\hbar} \right)^2 \quad (16)$$

where  $a_0 = \hbar^2/me^2$  is the 1st Bohr radius and  $\varepsilon = E/mc^2$ ;  $E$  is the Dirac energy:

$$E_{n,j} = mc^2 \left\{ 1 + Z^2\alpha^2 \left[ (n - j - 1/2) + \sqrt{(j + 1/2)^2 - Z^2\alpha^2} \right]^{-2} \right\}^{-1/2} \quad (17)$$

$\alpha = e^2/\hbar c$  is the fine structure constant and  $j = l \pm 1/2$  is the quantum number associated to the total angular momentum  $\vec{j} = \vec{l} + \vec{s}$ . The number  $\kappa$  is giving by the two relations  $\kappa = -(j + 1/2)$  if  $j = (l + 1/2)$  and  $\kappa = (j + 1/2)$  if  $j = (l - 1/2)$ .

We see that through  $\kappa$  in  $\langle r^{-2} \rangle$ , non-commutative corrections removes the degeneracy  $j = l + 1/2 = (l + 1) - 1/2$  and acts like the Lamb shift and the energy depends now on  $(n, j, l)$ , unlike the usual Dirac energies in (17). The equivalence between the two levels of  $l$  for the same  $j$  in (17) is accidental and is due to the Coulomb potential which is a special case. The additional term

in  $r^{-2}$  coming from non-commutativity breaks the symmetry and induces the differences found.

We consider in this study the effects of the finite mass of the nucleus, as we will discuss electronic and muonic atoms, where the reduced mass varies by a factor reaching 186 for hydrogen and 196 for the deuteron. To achieve this, we use the solution for two particles Dirac theory [40] and write the corrected Dirac energies as a shift of the usual ones:

$$E_{n,j} = mc^2 \left\{ 1 + \frac{1}{1+\eta} \left[ \frac{1}{sqr} - 1 \right] - \frac{\eta}{2(1+\eta)^3} \left[ \frac{1}{sqr} - 1 \right]^2 \right\} \quad (18)$$

Where  $\eta = m/M_N$  is the ratio between the orbiting particle mass  $m$  and the nucleus one  $M_N$  and  $sqr$  is the square expression in usual Dirac energies (17):

$$sqr = \sqrt{1 + Z^2\alpha^2 \left[ (n-j-1/2) + \sqrt{(j+1/2)^2 - Z^2\alpha^2} \right]^{-2}} \quad (19)$$

One retrieves (17) by putting  $\eta = 0$  in (18).

We use the shift (18) in the expression of  $\langle r^{-2} \rangle$  when computing the non-commutative corrections to the energies and thereby we generalize the recurrence relations of [39] for the case of a relativistic two-particle system; the result writes:

$$\Delta E_{n,j^\pm}^{(st)} = f^{(st)}(n, j^\pm, Z\alpha, \eta) (m^3 Z e^2 c^4 / \hbar^3) \theta_{st} \quad (20)$$

Here  $f^{(st)}(n, j^\pm, Z\alpha, \eta)$  is a dimensionless coefficient dependent on the parameters within the parentheses ( $j^\pm$  means  $j = l \pm 1/2$ ). To give an overview of this dependence, we develop its expression according to  $Z\alpha$ , and we limit ourselves to the 4th order:

$$f_{j^+}^{(st)} = \frac{-Z^2\alpha^2}{(1+\eta)^{3/2}j n^3} \left[ 1 + \left( \frac{6j^2+6j+1}{j(j+1)(2j+1)^2} + \frac{3}{(2j+1)n} - \frac{9\eta^2j+26\eta j+9\eta^2+24\eta+20j+18}{8(1+\eta)^2(j+1)n^2} \right) Z^2\alpha^2 \right] \quad (21a)$$

$$f_{j^-}^{(st)} = \frac{-Z^2\alpha^2}{(1+\eta)^{3/2}(j+1)n^3\hbar^3} \left[ 1 + \left( \frac{6j^2+6j+1}{j(j+1)(2j+1)^2} + \frac{3}{(2j+1)n} - \frac{9\eta^2j+26\eta j+2\eta+20j+2}{8(1+\eta)^2j n^2} \right) Z^2\alpha^2 \right] \quad (21b)$$

When putting  $\eta = 0$  and  $Z = 1$  in (20), (21a) and (21b), we recover the results of [31].

This concludes our study of the space-time case.

### 3.2 1st Order Corrections in Space-Space Non-Commutativity

For the space-space case and as mentioned before, the corrections are no longer central (13b) and we must use the full expression of the spinors  $\Psi(\vec{r})$  to perform the computations (we cite for example [41] and [42]). The method is very time consuming but the work has already been done in [28] and [30]. For the level

in which we are interested in this study ( $n = 2$ ), the corrections were found proportional to  $m^3 c^4 Z^4 \alpha^4 / \hbar^2$ ; we write their values from [28] ( $Z = 1$ ):

$$\Delta E_{2S_{1/2}}^{(ss)} = 0 \quad (22a)$$

$$\Delta E_{2P_{1/2}}^{(ss)} = \pm 6.75 \times 10^{19} \theta_{ss} eV / m^2 \quad (22b)$$

$$\Delta E_{2P_{3/2}}^{(ss)} = \pm (m_l + \frac{1}{2}) 6.75 \times 10^{19} \theta_{ss} eV / m^2 \quad (22c)$$

where  $\theta_{ss}$  is in  $m^2$  and  $m_l = 0, 1$ .

The sign  $\pm$  comes from the fact that  $\Delta H_{ss}$  removes the degeneracy according to the azimuthal quantum number  $m_l$  via the operator  $\vec{L}$ , and the  $1/2$  arises because the  $\Psi(\vec{r})$  for Dirac equation, are eigenstates of the operator  $\vec{J} = \vec{L} + \frac{1}{2} \text{diag}(\vec{\sigma}, \vec{\sigma})$  instead of  $\vec{L}$ .

We propose here to give another method to estimate the corrections in the space-space case. We start as done in [28] and [30], by writing:

$$\langle \Delta H_{ss} \rangle = -\frac{Ze^2}{4\hbar} \langle \Psi(r) | r^{-3} | \Psi(r) \rangle \langle \Psi(\vartheta, \varphi) | \vec{L} \cdot \vec{\theta}_{ss} | \Psi(\vartheta, \varphi) \rangle \quad (23)$$

The term in angular coordinates is a matrix, when diagonalized gives a contribution proportional to  $\theta_{ss} = \sqrt{\theta_j \theta^j}$ , and the proportion is a simple fraction noted  $a$ , so we approximate this term to  $a\theta_{ss}$ . We justify this by noting that when neglecting the fine-structure constant  $\alpha$  compared to unity, the  $\Psi(\vec{r})$  in Dirac equation becomes precisely the normalized Schrödinger eigenfunction provided we express the parameter  $\kappa$  in terms of  $l$  in the relativistic functions [41] and this gives the parameter  $\pm (m_l + \frac{1}{2})$  in  $\Delta E^{(ss)}$ .

For the radial term, and as done in space-time case, we use recurrence relation from [39]:

$$\left\langle \frac{1}{r^3} \right\rangle = \frac{2 [3\kappa^2 \varepsilon^2 - 3\kappa \varepsilon - (\kappa^2 - Z^2 \alpha^2) + 1] (1 - \varepsilon^2)^{3/2}}{\sqrt{\kappa^2 - Z^2 \alpha^2} [(\kappa^2 - Z^2 \alpha^2) - 1] [4(\kappa^2 - Z^2 \alpha^2) - 1]} \left( \frac{mc}{\hbar} \right)^3 \quad (24)$$

with the same notations as in (16).

In this case too, we consider the effects of the finite mass of the nucleon. When using the shift (18) in the expression of  $\langle r^{-3} \rangle$ , the non-commutative corrections to the energies are:

$$\Delta E_{n,j^\pm}^{(ss)} = f^{(ss)}(n, j^\pm, Z\alpha, \eta) (m^3 Z e^2 c^3 / \hbar^3) a \theta_{st} \quad (25)$$

with the same notations as in space-time case. The approximate solution of the coefficient  $f$  is:

$$f_{j^+}^{(ss)} = \frac{(1+\eta)^{-3/2} Z^3 \alpha^3}{j(2j-1)(2j+1)n^3} \left[ 1 + \left( \frac{48j^2 + 36j^2 - 8j - 3}{j(2j-1)(2j+1)^2(2j+3)} + \frac{3}{(2j+1)n} - \frac{3 \cdot 6\eta^2 j + 20\eta j + 9\eta^2 + 22\eta + 16j + 16}{8(1+\eta)^2(2j+3)n^2} \right) Z^2 \alpha^2 \right] \quad (26a)$$

$$f_{j^-}^{(ss)} = \frac{(1+\eta)^{-3/2} Z^3 \alpha^3}{(j+1)(2j+1)(2j+3)n^3} \left[ 1 + \left( \frac{48j^2 + 108j^2 + 64j + 7}{(j+1)(2j-1)(2j+1)^2(2j+3)} + \frac{3}{(2j+1)n} - \frac{3 \cdot 6\eta^2 j + 20\eta j - 3\eta^2 - 2\eta + 16j}{8(1+\eta)^2(2j-1)n^2} \right) Z^2 \alpha^2 \right] \quad (26b)$$



To justify our approach, we apply it to the level  $n = 2$  of the hydrogen and we find ( $Z = 1$ ):

$$\Delta E_{2S_{1/2}}^{(ss)} = \pm 9.11531 \times 10^{-4} (m^3 e^2 c^3 / \hbar^3) a \theta_{ss} \quad (27a)$$

$$\Delta E_{2P_{1/2}}^{(ss)} = \pm 5.05790 \times 10^{-9} (m^3 e^2 c^3 / \hbar^3) a \theta_{ss} \quad (27b)$$

$$\Delta E_{2P_{3/2}}^{(ss)} = \pm 4.04467 \times 10^{-9} (m^3 e^2 c^3 / \hbar^3) a \theta_{ss} \quad (27c)$$

For the  $2S_{1/2}$  state, we put  $a = 0$  because the  $\vec{L}$  operator vanishes in this state and we retrieve the result of [28] here. For the  $P$ -states, comparing our results to those coming from [28] gives:

$$a(2P_{1/2}) = 0.53 \ \& \ a(2P_{3/2}) = 0.33 - 1.00 \quad (28)$$

These values show that our method gives results very close to the exact values from [28] and [30] but it is much less tedious. It also has the advantage of providing a general formulation of the results, which is not the case of the pre-cited studies.

## 4 Non-Commutativity in H-Like Atoms

Now we apply the expressions of non-commutative corrections found in the previous section on hydrogen and deuterium whether electronic or muonic, with the aim to find an explanation for the different results of aforementioned experiments. We will start with the result that has generated the more debates and that relates to the muonic hydrogen. Then we will discuss some aspects concerning the radii of proton and deuteron and atomic spectroscopy through muonic deuterium, electronic hydrogen and electronic deuterium.

### 4.1 Muon Hydrogen and Non-Commutativity

In this section, we will apply our study of both space-time and space-space cases of non-commutativity to the muon hydrogen and especially to the transitions used to compute the charge radius of the proton (1a) and (1b). Using the value of the charge radius given by CODATA 2010  $0.8775(51)fm$  [2], the results obtained for the precedent transitions differ from those found in experiments on muonic hydrogen [1], by an amount equal to  $0.32meV$ . Although the difference is very small, it is significant given the precision of the experiments used.

We compute the non-commutative corrections to these transitions for the space-time case using (16) (17) (18) and (19); we find :

$$\Delta E_{2P_{1/2}}^{(st)} - \Delta E_{2S_{1/2}}^{(st)} = 6.51015 \times 10^{16} (\theta_{st} eV^2) eV \quad (29a)$$

$$\Delta E_{2P_{3/2}}^{(st)} - \Delta E_{2S_{1/2}}^{(st)} = 6.51044 \times 10^{16} (\theta_{st} eV^2) eV \quad (29b)$$

For the space-space type, we use (19) (23) (24) and (25):

$$\Delta E_{2P_{1/2}}^{(ss)} - \Delta E_{2S_{1/2}}^{(ss)} = 3.93686 \times 10^{13} (\theta_{ss} eV^2) eV \quad (30a)$$

$$\Delta E_{2P_{3/2}}^{(ss)} - \Delta E_{2S_{1/2}}^{(ss)} = 2.96894 \times 10^{13} (\theta_{ss} eV^2) eV \quad (30b)$$

We have chosen  $a(2S_{1/2}) = 0$ ,  $a(2P_{1/2}) = 0.5$  &  $a(2P_{3/2}) = 1$  to get back the results of [28].

Comparing these results to the deviation  $0.32 \times 10^{-3} eV$ , we compute the values of the parameter of non-commutativity that is required to fill the gap:

$$\Delta E_{2P_{1/2}}^{(st)} - \Delta E_{2S_{1/2}}^{(st)} = 0.32 meV \implies \theta_{st} = (14.286 GeV)^{-2} \quad (31a)$$

$$\Delta E_{2P_{3/2}}^{(st)} - \Delta E_{2S_{1/2}}^{(st)} = 0.32 meV \implies \theta_{st} = (14.286 GeV)^{-2} \quad (31b)$$

$$\Delta E_{2P_{1/2}}^{(ss)} - \Delta E_{2S_{1/2}}^{(ss)} = 0.32 meV \implies \theta_{ss} = (0.305 GeV)^{-2} \quad (31c)$$

$$\Delta E_{2P_{3/2}}^{(ss)} - \Delta E_{2S_{1/2}}^{(ss)} = 0.32 meV \implies \theta_{ss} = (0.351 GeV)^{-2} \quad (31d)$$

The value obtained in the case of space-space non-commutativity  $\theta_{ss} \approx (0.3 GeV)^{-2}$  exceeds the limit obtained for this parameter from studies on  $eH$  Lamb shift  $\theta_{ss} \leq (0.6 GeV)^{-2}$  [33]. If we use this limit to compute the radius, we find  $0.86409 fm$  which is outside the experimental limits for  $r_{\mu H}$  [1] and [3]; So this effect is ruled out.

For the case of the space-time non-commutativity, the value  $\theta_{st} \approx (14.3 GeV)^{-2}$  is obtained and it is in agreement with the limit  $\theta_{st} \leq (6 GeV)^{-2}$  determined by Lamb shift spectroscopy in  $eH$  [33]. It should also be noted that the corrections are the same for both levels  $2P_{1/2}$  and  $2P_{3/2}$ ; this is in agreement with the two relations (1a) and (1b), where the terms in  $r$  are equal for both transitions.

This ends our quantitative analysis of non-commutative effects on the spectrum of muonic hydrogen and on the charge radius of the proton.

## 4.2 Lamb Shift in Muon Deuterium

Now, we study the consequences of the noncommutative correction found for  $\mu H$  on the muon deuterium  $\mu D$ . The  $2P - 2S$  Lamb shift has been extensively studied to see if the results obtained with  $\mu H$  are confirmed by this new muon system. The theoretical value of this transition depends on the deuteron radius and is given by the formula [43]:

$$\frac{\Delta(2P_{1/2} \rightarrow 2S_{1/2})}{meV} = 230.2972(400) - 6.10940 \frac{r^2}{fm^2} + 0.0448 \frac{r^3}{fm^3} \quad (32a)$$

The Mainz collaboration has studied the Lamb-shift of  $\mu D$  and found that the experimental value differs from the theoretical one by  $0.383 meV$  [44] (and references therein). Although the difference is only  $3\sigma$ , there is scope for study in this phenomenon. Several studies have looked at this problem and especially on the possible correction to the two-photon exchange (proportional to  $r^3$ ) to

find a solution, but without conclusive results until now [45] [46] (For a review one can see [47]).

We propose to solve this problem by using the same procedure used to  $\mu H$ . To do this, we calculate the non-commutative correction to the transition for space-time case. From (20), (21a) and (21b), the non-commutative correction to the Lamb shift in hydrogen-like atoms is given with the general expression:

$$\begin{aligned} \Delta E_{n,j}^{(st)}(Lamb\ shift) &= \Delta E_{n,j=(l+1)-1/2}^{(st)} - \Delta E_{n,j=l+1/2}^{(st)} \\ &= \frac{1}{(1+\eta)^{3/2}} \frac{m^3 c^4 e^2 Z^3 \alpha^2 \theta_{st}}{j(j+1)n^3 \hbar^3} \left[ 1 + \left( \frac{6j^2+6j+1}{j(j+1)(2j+1)^2} + \frac{3}{(2j+1)n} - \frac{9\eta^2+22\eta+16}{8(1+\eta)^2 n^2} \right) Z^2 \alpha^2 \right] \end{aligned} \quad (33)$$

For the  $2P_{1/2} \rightarrow 2S_{1/2}$  case in  $\mu D$ , we take  $\eta = m_\mu/M_D$ ,  $n = 2$  and  $j = 1/2$ ; we find:

$$\Delta E_{2,1/2}^{(st)}(LS) = 7.03726 \times 10^{16} (\theta_{st} eV^2) eV \quad (34)$$

The same value is obtained when using the exact expressions (20). Putting the value  $\theta_{st} \approx (14.3eV)^{-2}$  (31a) obtained from  $\mu H$  in (34), we get  $\Delta E^{(st)} = 0.348meV$ ; this value is approximately equal to the discrepancy between theoretical and experimental results  $0.383meV$ . We see that the same parameter corrects the two observed discrepancies in both  $\mu H$  and  $\mu D$ .

Be noted that the value  $0.348meV$  in  $\mu D$  is very close the value  $0.32meV$  that counts for  $\mu H$  and this is easily explained by the fact that the difference between the two systems comes from the factor  $m^3 (1+\eta)^{-3/2}$  in (33) and the ratio of the numerical quantities is  $0.93 \approx 1$ .

### 4.3 $2S-1S$ Transition in Hydrogen and Deuterium

It should be noted that the difference between the radii of proton and deuteron is a very well-defined parameter using the  $2S - 1S$  transition which is one of the most accurate measured quantities [48]. This transition is used because the effects due to nuclear size ( $ns$ ) are, to the first order, nonzero only for these states [34]:

$$\Delta E_{ns} = \frac{2}{3} \frac{\mu^3 c^4 Z^4 \alpha^4}{\hbar^2 n^3} \langle r^2 \rangle \delta_{l0} \quad (35)$$

We see that knowing the reduced masses  $\mu$  allows us to accurately evaluate radii of nuclei from high precision spectroscopy.

The most recent measurement of the hydrogen-deuterium isotope shift is [49]:

$$\Delta f_{ex}(H - D) = f_H^{1S-2S} - f_D^{1S-2S} = 670994334606(15)Hz \quad (36)$$

and the most recent theoretical evaluation is [50]:

$$\Delta f_{th}(H - D) = f_H^{1S-2S} - f_D^{1S-2S} = 670994346(23)kHz \quad (37)$$

Both values agree well within the limits given.

We will evaluate the contribution coming from space-time non-commutativity to this shift. We use the relations from (15) to (20), to compute the non-commutative corrections to the transition in both hydrogen and deuterium:

$$\Delta E_{2S}^{(st)}(H) - \Delta E_{1S}^{(st)}(H) = hf_H^{1S-2S} = 9.06690 \times 10^{10} (\theta_{st} eV^2) eV \quad (38a)$$

$$\Delta E_{2S}^{(st)}(D) - \Delta E_{1S}^{(st)}(D) = hf_D^{1S-2S} = 9.07060 \times 10^{10} (\theta_{st} eV^2) eV \quad (38b)$$

Inserting the value of the parameter  $\theta_{st}$  found for  $\mu H$  (31a) in these expressions, we find the non-commutative correction to the hydrogen-deuterium isotope shift:

$$\Delta f_{th}^{(nc)}(H - D) = f_H^{1S-2S} - f_D^{1S-2S} = 275.488 Hz \quad (39)$$

This contribution don't fill the gap of  $12 kHz$  between  $\Delta f_{th}$  and  $\Delta f_{ex}$ , but it improves slightly the agreement between the two values. This confirms the fact that there are no doubts about the results of experiments on  $eH$  spectroscopy because they are so accurate and one has to look the side of muonic systems.

## 5 Conclusion

In this work, we studied the corrections induced by a non-commutative structure of space-time, in its two versions space-space and space-time, on the spectrum of hydrogen-like atoms. We have applied our study to the muonic hydrogen and this with the aim to solve the puzzle of proton radius, because we think that the experiments used to study this phenomenon are so developed that we can not doubt of their results, and therefore one must look on the side of theory of atomic spectroscopy used to compute the radius.

In this study, we considered the effects of the mass of the nucleus of the non-commutative corrections and thus we have improved previous works in this area. This allowed us to consider the difference caused by changing the nucleus (from proton to deuteron) in addition to that which occurs when changing the orbiting particle (from electron to muon).

It should be noted that the effects of the nucleus shape on the energy levels of the atom are proportional to the third power of the mass of the orbiting particle; this is easily understood by the fact that the Bohr radius ( $a_0 = \hbar^2/me^2$ ) is inversely proportional to the mass and thus the particle is that much nearer the nucleus, that its mass is greater; and this makes it more sensitive to these effects.

We have demonstrated that the effects of non-commutativity are also proportional to the third power of the mass of the particle because it distorts the Coulomb potential and adds a term proportional to  $r^{-2}$  in space-time case and to  $r^{-3}$  in space-space case. It is for this reason also that the effects decreases with increasing values of quantum numbers as can be seen in the different relations of the spectrum corrections (because the term  $r^{-n}$  with  $n > 1$  is very steep for small values of  $r$ ). This is why we use this theory to explain the puzzle

because its effects are different depending on whether it is applied to muonic hydrogen or electronic hydrogen. The shifts in the spectrum are more noticeable in muon  $H$  than in ordinary ones, and this explains the fact that experiments on  $\mu H$  spectroscopy give results that are different from those obtained with  $eH$ .

In the case of space-space non-commutativity, the parameter required to resolve the puzzle is  $\theta_{ss} \approx (0.35 GeV)^{-2}$ . This value exceeds the limit obtained for this parameter from studies on  $eH$  Lamb shift  $\theta_{ss} \leq (0.6 GeV)^{-2}$  [33]. If we use this limit to compute the radius, we find  $0.86409 fm$  which is outside the experimental limits for  $r_{\mu H}$  [1] and [3]. Another problem arises with in this case; it is the  $\pm$  sign in the corrections (due to the presence of the azimuthal quantum number in their expressions). This sign means that the corrected value of the radius ranges from  $0.84169 fm$  to  $0.91192 fm$  in violation of  $\mu H$  experiences.

In the case of the space-time non-commutativity, the value  $\theta_{st} \approx (14.3 GeV)^{-2}$  has been obtained and it is in agreement with the limit determined by Lamb shift spectroscopy in  $eH$   $\theta_{st} \leq (6 GeV)^{-2}$  [33]. It was also found in this case, that the corrections are the same for both levels  $2P_{1/2}$  and  $2P_{3/2}$  although we found that corrections remove the degeneracy of the Dirac energies with respect to the total angular momentum quantum number  $j = l + 1/2 = (l + 1) - 1/2$  (non-commutativity acts like the Lamb Shift here). This is in agreement with the two relations (27a) and (27b), where the terms in  $r$  are equal for both transitions. This is not true for the space-space case because the corrections of the two levels differ from one another (29a) (29b). We say that this is a consequence of the fact that the correction term to the Coulomb potential is proportional to  $r^{-2}$  in the space-time case, and so as we have previously mentioned in [31], we assimilate it to the field of a central dipole. In other words, the action of space-time non-commutativity is equivalent to consider the extended charged nature of the proton in the nucleus, which is the principal characteristic studied in  $\mu H$  experiments.

When applying the result obtained from the study of  $\mu H$  in  $\mu D$ , we found a correction of  $0.348 meV$  which is almost exactly equal to the difference between theory and experiment for this system. The very close values of the corrections in these two systems  $\mu H$  and  $\mu D$  are easily explained by the fact that the ratio between the reduced masses, of the two is  $0.95 \approx 1$  (The disagreements between theory and experiment in both systems are almost equal).

Using the same result coming from  $\mu H$  for the  $eH$ - $eD$  isotope shift, the correction found improves the agreement between theoretical and experimental results (which was already excellent). The same reasoning as above is used, and we say that the corrections in electronic systems either  $eH$  or  $eD$  are infinitely small compared to those of muonic systems; the ratio between the reduced masses of the two is  $\approx 10^{-7}$ .

Eyes are now turned to the results of experimental on  $\mu p$  scattering and  $\mu He$  spectroscopy to see whether the phenomenon is spectroscopic or is it due to the nature of the particles. On the side of electronic systems, there is practically no doubt on their veracity; the radius of the proton was even calculated in a model independent way from  $ep$  scattering [51] [52] and the results confirm CODATA

value.

It was reported in our work [31], that the limit on the parameter  $\theta_{st} \approx (1\text{TeV})^{-2}$ , and so our result here is greater than the latter; however it was pointed out to us that such a limit ( $\theta_{st} \approx (1\text{TeV})^{-2}$ ) should not only be estimated according to experimental precision calculations but rather on the disagreement between experiment and theory, which requires a correction of the limit of [31] and this is what is done in this work for the muon hydrogen.

We can also evoke that the proton raises other questions about its properties in addition to the one discussed in this article and we can mention as an example the origin of its spin or what is called "spin crisis" in [53] or "proton spin puzzle" in [54].

## References

- [1] R. Pohl et al., *The size of the proton*, Nature **466** pp 213-216 (2010)
- [2] P. J. Mohr, B. N. Taylor and D. B. Newell, *CODATA Recommended Values of the Fundamental Physical Constants: 2010*, Rev. Mod. Phys. **84**, pp 1527-1605 (2012)
- [3] A. Antognini et al., *Proton Structure from the Measurement of 2S-2P Transition Frequencies of Muonic Hydrogen*, Science **339** pp. 417-420 (2013)
- [4] J. D. Carroll, A. W. Thomas, J. Rafelski and G. A. Miller, *Nonperturbative relativistic calculation of the muonic hydrogen spectrum*, Phys. Rev. **A 84**, 012506 (2011)
- [5] P. Indelicato, *Nonperturbative evaluation of some QED contributions to the muonic hydrogen n=2 Lamb shift and hyperfine structure*, Phys. Rev. **A 87**, 022501 (2013)
- [6] R. Pohl, R. Gilman, G. A. Miller and K. Pachucki, *Muonic hydrogen and the proton radius puzzle*, Annu. Rev. Nucl. Part. Sci **63** pp 175-204 (2013)
- [7] A. Antognini et al., *Theory of the 2S-2P Lamb shift and 2S hyperfine splitting in muonic hydrogen*, Ann. Phys. **331** pp 127-145 (2013)
- [8] I. C. Cloët and G. A. Miller, *Third Zemach moment of the proton*, Phys. Rev. **C 83**, 012201(R) (2011)
- [9] J. P. Karr and L. Hillico, *Why Three-Body Physics Does Not Solve the Proton-Radius Puzzle*, Phys. Rev. Lett. **109**, 103401 (2012)
- [10] V. Barger, C. W. Chiang, W. Y. Keung and D. Marfatia, *Proton Size Anomaly*, Phys. Rev. Lett. **106**, 153001 (2011)
- [11] G. A. Miller, A. W. Thomas, J. D. Carroll, and J. Rafelski, *Toward a resolution of the proton size puzzle*, Phys. Rev. **A 84**, 020101(R) (2011)

- [12] G. A. Miller, *Proton polarizability contribution: Muonic hydrogen Lamb shift and elastic scattering*, Phys. Lett **B 718** pp 1078-1082 (2013)
- [13] U. D. Jentschura, *Lamb shift in muonic hydrogen-I. Verification and update of theoretical predictions*, Ann. Phys. **326** pp 500–515 (2011)
- [14] U. D. Jentschura, *Lamb shift in muonic hydrogen-II. Analysis of the discrepancy of theory and experiment*, Ann. Phys. **326** pp 516–533 (2011)
- [15] Gilman et al., Technical Design Report Paul Scherrer Institute Experiment R-12-01.1 (2013) arXiv:1303.2160
- [16] A. Connes, *Noncommutative Geometry*, (Academic Press, CA, 1994)
- [17] N. Seiberg and E. Witten, *String theory and Noncommutative Geometry*, JHEP **09**, 032 (1999)
- [18] J. Bellissard, H. Schulz-Baldes and A. Van Elst, *The Non Commutative Geometry of the Quantum Hall Effect*, J. Math. Phys. **35** pp 5373-5471 (1994)
- [19] J. Bellissard, *Noncommutative Geometry of Aperiodic Solids*, in *Proc. of the Summer School Geometric and Topological Methods for Quantum Field Theory*, eds., H. Ocampo, S. Paycha and A. Cardona (World Sci. Publishing, River Edge, NJ, 2003) pp. 86-156.
- [20] A. P. Polychronakos, *Non-Commutative Fluids*, Prog. Math. Phys. **53** pp 109-159 (2007) (and the references therein)
- [21] A. Connes and M. Marcolli, *From Physics to Number Theory via Noncommutative Geometry*, Frontiers in Numbers Theory, Physics, and Geometry I, Part II, pp 269-349 (Springer, 2006)
- [22] A. Sitarz, *Non-Commutative Geometry and the Ising Model*, J. Phys. A: Math. Gen. **26** pp 5305-5312 (1993)
- [23] A. Bérard and H. Mohrbach, *Spin Hall Effect and Berry Phase of Spinning Particles*, Phys. Lett. **A352** pp 190-195(2006)
- [24] D. B. Tweed, T. P. HalsWanter, V. Happe and M. Fetter, *Non-Commutativity in the Brain*, Nature **399** pp 261-263 (1999)
- [25] M. R. Douglas and N. A. Nekrasov, *Noncommutative Field Theory*, Rev. Mod. Phys. **73** pp 977-1029 (2001)
- [26] R. J. Szabo, *Quantum Field Theory on Noncommutative Spaces*, Phys. Rept. **378**, pp 207-299 (2003)
- [27] M. Chaichian, M. M. Sheikh-Jabbari and A. Tureanu, *Hydrogen Atom Spectrum and the Lamb Shift in Noncommutative QED*, Phys. Rev. Lett. **86**, 2716 (2001)

- [28] T. C. Adorno, M. C. Baldiotti, M. Chaichian, D. M. Gitman and A. Tureanu, *Dirac Equation in Noncommutative Space for Hydrogen Atom*, Phys. Lett. **B682**, pp 235-239 (2009)
- [29] W. O. Santos and A. M. C. Souza, *The Anomalous Zeeman Effect for the Hydrogen Atom in Noncommutative Space*, Int. J. Theor. Phys. **51** pp 3882-3890 (2011)
- [30] L. Khodja and S. Zaim, *New treatment of the noncommutative Dirac equation with a Coulomb potential*, Int. J. Mod. Phys. **A 27**, 1250100 (2012)
- [31] M. Moumni, A. BenSlama and S. Zaim, *Spectrum of Hydrogen Atom in Space-Time Non-Commutativity*, Afr. Rev. Phys. **07** pp 83-94 (2012)
- [32] M. Moumni, A. BenSlama and S. Zaim, *Relativistic spectrum of hydrogen atom in the space-time non-commutativity*, in *Proc 8th Int Conf on Progress in Theoretical Physics* AIP Conf. Proc., Vol 1444, eds. Mebarki et al. (2012) pp 253-257
- [33] A. Stern, *Particlelike Solutions to Classical Noncommutative Gauge Theory*, Phys. Rev. **D78**, 065006 (2008)
- [34] J. L. Friar, *Nuclear finite-size effects in light muonic atoms*, Ann. Phys. **122**, pp 151-196 (1979)
- [35] M. I. Eides, H. Grotch and V. A. Shelyuto, *Theory of light hydrogenlike atoms*, Phys. Rep. **342** pp 63–261 (2001)
- [36] S. Dulat and K. Li, *The Aharonov-Casher Effect for Spin-1 Particles in Non-Commutative Quantum Mechanics*, Eur. Phys.J. **C54**, pp 333-337 (2008)
- [37] S. Fabi, B. Harms and A. Stern, *Noncommutative Corrections to the Robertson-Walker Metric*, Phys. Rev. **D78**, 065037 (2008)
- [38] M. Chaichian, A. Tureanu, M. R. Setare and G. Zet, *On Black Holes and Cosmological Constant in Noncommutative Gauge Theory of Gravity*, JHEP **0804**, 064 (2008)
- [39] S. K. Suslov and B. Trey, *The Hahn Polynomials in the Nonrelativistic and Relativistic Coulomb Problems*, J. Math. Phys. **49**, 012104 (2008)
- [40] J. Sapirstein, *Quantum Electrodynamics*, in *Atomic, Molecular, & Optical Physics Handbook*, edited by G. W. F. Drake, (AIP Press, New York 1996), pp. 327–340
- [41] H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One- and Two-Electron Atoms*, (Academic Press, NY 1957)
- [42] W. Greiner, *Relativistic Quantum Mechanics Wave Equations*, 3rd edn (Springer, Berlin, 2000)



- [43] E. Borie, *Lamb shift in light muonic atoms - Revisited*, Ann. Phys. **327**, pp 733-763 (2012)
- [44] M. Gorshteyn, *Nuclear and Hadronic Contributions to Lamb Shift In Muonic Deuterium*, Talk given in *Workshop to Explore Physics Opportunities with Intense, Polarized Electron Beams up to 300 MeV*, MIT, Cambridge, Massachusetts, March 14-16, 2013
- [45] K. Pachucki, *Nuclear Structure Corrections in Muonic Deuterium*, Phys. Rev. Lett. **106**, 193007 (2011)
- [46] A. A. Krutov and A. P. Martynenko, *Lamb shift in the muonic deuterium atom*, Phys. Rev. **A 84**, 052514 (2011)
- [47] R. Pohl, *Muonic News*, Talk given at *ECT Workshop on the "Proton Radius Puzzle"*, Trento, Italy, Oct 29 to Nov 2 (2012)
- [48] C. G. Parthey et al., *Improved Measurement of the Hydrogen  $1S - 2S$  Transition Frequency*, Phys. Rev. Lett. **107**, 203001 (2011)
- [49] C. G. Parthey et al.; *Precision Measurement of the Hydrogen-Deuterium  $1S - 2S$  Isotope Shift*, Phys. Rev. Lett. **104**, 233001 (2010)
- [50] U. D. Jentschura et al., *Hydrogen-deuterium isotope shift: From the  $1S - 2S$  transition frequency to the proton-deuteron charge-radius difference*, Phys. Rev. **A 83**, 042505 (2011)
- [51] R. J. Hill and G. Paz, *Model-independent extraction of the proton charge radius from electron scattering*, Phys. Rev. **D 82**, 113005 (2010)
- [52] I. Sick, *Problems with proton radii*, Prog. Partic. Nucl. Phys. **67** pp 473-478 (2012)
- [53] A. W. Thomas, A. Casey and H. H. Matevosyan, *What we Know and Don't Know about the Origin of the Spin of the Proton*, Int. J. Mod. Phys. **A 25**, pp. 4149-4162 (2010)
- [54] C. A. Aidala, S. D. Bass, D. Hasch and G. K. Mallot, *The spin structure of the nucleon*, Rev. Mod. Phys. **85**, pp. 655-691 (2013)